

WHAT IS CLAIMED IS:

1. Hybrid maize seed designated X1069G (commercial designation), representative seed of said hybrid X1069G having been deposited under ATCC accession number \_\_\_\_\_.
2. A maize plant, or its parts, produced by the seed of claim 1.
3. Pollen of the plant of claim 2.
4. An ovule of the plant of claim 2.
5. A tissue culture of regenerable cells of a hybrid maize plant X1069G (commercial designation), representative seed of said hybrid maize plant X1069G having been deposited under ATCC accession number \_\_\_\_\_, wherein the tissue regenerates plants capable of expressing all the morphological and physiological characteristics of said hybrid maize plant X1069G.
6. A tissue culture according to claim 5, the cells or protoplasts being from a tissue selected from the group consisting of leaves, pollen, embryos, roots, root tips, anthers, silks, flowers, kernels, ears, cobs, husks, and stalks.
7. A maize plant, or its parts, regenerated from the tissue culture of claim 5 and capable of expressing all the morphological and physiological characteristics of hybrid maize plant X1069G (commercial designation), representative seed having been deposited under ATCC accession number \_\_\_\_\_.
8. The maize plant of claim 2 wherein said plant is male sterile.
9. A method for developing a maize plant in a maize plant breeding program using plant breeding techniques, which include employing a maize plant, or its parts, as a source of plant breeding material, comprising: obtaining the maize plant, or its parts, of claim 2 as a source of said breeding material.
10. The maize plant breeding program of claim 9 wherein plant breeding techniques are selected from the group consisting of: recurrent selection, backcrossing,

pedigree breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, and transformation.

11. A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 2, said maize plant capable of expressing a combination of at least two X1069G (commercial designation) traits selected from the group consisting of: excellent yield potential, good stalk lodging resistance, good root lodging resistance, good early growth, good stay green, good test weight, very good dry down, very good ear retention, very good husk cover, dependable drought stress tolerance, moderate resistance to Gray Leaf Spot, moderate resistance to Northern Leaf Blight, moderate resistance to Eye Spot, moderate resistance to Fusarium Ear Rot, moderate resistance to Gibberella Ear Rot, moderate resistance to Common Rust, excellent resistance to head smut, moderate resistance to European Corn Borer first and second generation, suited to the Central Corn Belt, Northwest, Northcentral, Northeastern, and Western regions of the United States, and a relative maturity of approximately 105 (106 for physiological maturity) based on the Comparative Relative Maturity Rating System for harvest moisture of grain.
12. A hybrid maize plant according to claim 2, wherein the genetic material of said plant contains one or more transgenes.
13. A method for developing a maize plant in a maize plant breeding program using plant breeding techniques, which include employing a maize plant, or its parts, as a source of plant breeding material, comprising: obtaining the maize plant, or its parts, of claim 12 as a source of said breeding material.
14. The maize plant breeding program of claim 13 wherein plant breeding techniques are selected from the group consisting of: recurrent selection, backcrossing, pedigree breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, and transformation.
15. A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 12, said maize plant capable of expressing a combination of at least two X1069G (commercial designation) traits selected from the group consisting of: excellent yield potential, good stalk lodging resistance,

good root lodging resistance, good early growth, good stay green, good test weight, very good dry down, very good ear retention, very good husk cover, dependable drought stress tolerance, moderate resistance to Gray Leaf Spot, moderate resistance to Northern Leaf Blight, moderate resistance to Eye Spot, moderate resistance to Fusarium Ear Rot, moderate resistance to Gibberella Ear Rot, moderate resistance to Common Rust, excellent resistance to head smut, moderate resistance to European Corn Borer first and second generation, suited to the Central Corn Belt, Northwest, Northcentral, Northeastern, and Western regions of the United States, and a relative maturity of approximately 105 (106 for physiological maturity) based on the Comparative Relative Maturity Rating System for harvest moisture of grain.

16. A hybrid maize plant according to claim 2, wherein the genetic material of said plant contains one or more genes transferred by backcrossing.
17. A method for developing a maize plant in a maize plant breeding program using plant breeding techniques, which include employing a maize plant, or its parts, as a source of plant breeding material, comprising: obtaining the maize plant, or its parts, of claim 16 as a source of said breeding material.
18. The maize plant breeding program of claim 17 wherein plant breeding techniques are selected from the group consisting of: recurrent selection, backcrossing, pedigree breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, and transformation.
19. A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 16, said maize plant capable of expressing a combination of at least two X1069G (commercial designation) traits selected from the group consisting of: excellent yield potential, good stalk lodging resistance, good root lodging resistance, good early growth, good stay green, good test weight, very good dry down, very good ear retention, very good husk cover, dependable drought stress tolerance, moderate resistance to Gray Leaf Spot, moderate resistance to Northern Leaf Blight, moderate resistance to Eye Spot, moderate resistance to Fusarium Ear Rot, moderate resistance to Gibberella Ear Rot, moderate resistance to Common Rust, excellent resistance to head smut,

moderate resistance to European Corn Borer first and second generation, suited to the Central Corn Belt, Northwest, Northcentral, Northeastern, and Western regions of the United States, and a relative maturity of approximately 105 (106 for physiological maturity) based on the Comparative Relative Maturity Rating System for harvest moisture of grain.

20. A maize plant, or its parts, having all the morphological and physiological characteristics of the plant of claim 2.
21. The maize plant of claim 20 wherein said maize plant is male sterile.
22. A method for developing a maize plant in a maize plant breeding program using plant breeding techniques, which include employing a maize plant, or its parts, as a source of plant breeding material, comprising: obtaining the maize plant, or its parts, of claim 20 as a source of said breeding material.
23. The maize plant breeding program of claim 22 wherein plant breeding techniques are selected from the group consisting of: recurrent selection, backcrossing, pedigree breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, and transformation.
24. A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 20, said maize plant capable of expressing a combination of at least two X1069G (commercial designation) traits selected from the group consisting of: excellent yield potential, good stalk lodging resistance, good root lodging resistance, good early growth, good stay green, good test weight, very good dry down, very good ear retention, very good husk cover, dependable drought stress tolerance, moderate resistance to Gray Leaf Spot, moderate resistance to Northern Leaf Blight, moderate resistance to Eye Spot, moderate resistance to Fusarium Ear Rot, moderate resistance to Gibberella Ear Rot, moderate resistance to Common Rust, excellent resistance to head smut, moderate resistance to European Corn Borer first and second generation, suited to the Central Corn Belt, Northwest, Northcentral, Northeastern, and Western regions of the United States, and a relative maturity of approximately 105 (106 for

physiological maturity) based on the Comparative Relative Maturity Rating System for harvest moisture of grain.

25. A hybrid maize plant according to claim 20, wherein the genetic material of said plant contains one or more transgenes.
26. A method for developing a maize plant in a maize plant breeding program using plant breeding techniques, which include employing a maize plant, or its parts, as a source of plant breeding material, comprising: obtaining the maize plant, or its parts, of claim 25 as a source of said breeding material.
27. The maize plant breeding program of claim 26 wherein plant breeding techniques are selected from the group consisting of: recurrent selection, backcrossing, pedigree breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, and transformation.
28. A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 25, said maize plant capable of expressing a combination of at least two X1069G (commercial designation) traits selected from the group consisting of: excellent yield potential, good stalk lodging resistance, good root lodging resistance, good early growth, good stay green, good test weight, very good dry down, very good ear retention, very good husk cover, dependable drought stress tolerance, moderate resistance to Gray Leaf Spot, moderate resistance to Northern Leaf Blight, moderate resistance to Eye Spot, moderate resistance to Fusarium Ear Rot, moderate resistance to Gibberella Ear Rot, moderate resistance to Common Rust, excellent resistance to head smut, moderate resistance to European Corn Borer first and second generation, suited to the Central Corn Belt, Northwest, Northcentral, Northeastern, and Western regions of the United States, and a relative maturity of approximately 105 (106 for physiological maturity) based on the Comparative Relative Maturity Rating System for harvest moisture of grain.
29. A hybrid maize plant according to claim 20, wherein the genetic material of said plant contains one or more genes transferred by backcrossing.

30. A method for developing a maize plant in a maize plant breeding program using plant breeding techniques, which include employing a maize plant, or its parts, as a source of plant breeding material, comprising: obtaining the maize plant, or its parts, of claim 29 as a source of said breeding material.
31. The maize plant breeding program of claim 30 wherein plant breeding techniques are selected from the group consisting of: recurrent selection, backcrossing, pedigree breeding, restriction fragment length polymorphism enhanced selection, genetic marker enhanced selection, and transformation.
32. A maize plant, or its parts, wherein at least one ancestor of said maize plant is the maize plant, or its parts, of claim 29, said maize plant capable of expressing a combination of at least two X1069G (commercial designation) traits selected from the group consisting of: excellent yield potential, good stalk lodging resistance, good root lodging resistance, good early growth, good stay green, good test weight, very good dry down, very good ear retention, very good husk cover, dependable drought stress tolerance, moderate resistance to Gray Leaf Spot, moderate resistance to Northern Leaf Blight, moderate resistance to Eye Spot, moderate resistance to Fusarium Ear Rot, moderate resistance to Gibberella Ear Rot, moderate resistance to Common Rust, excellent resistance to head smut, moderate resistance to European Corn Borer first and second generation, suited to the Central Corn Belt, Northwest, Northcentral, Northeastern, and Western regions of the United States, and a relative maturity of approximately 105 (106 for physiological maturity) based on the Comparative Relative Maturity Rating System for harvest moisture of grain.